

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/96, A23L 1/00, C12N 9/16, 9/24, 9/42	A1	(11) International Publication Number: WO 97/05245 (43) International Publication Date: 13 February 1997 (13.02.97)
(21) International Application Number: PCT/EP96/03286 (22) International Filing Date: 24 July 1996 (24.07.96) (30) Priority Data: 95202066.7 28 July 1995 (28.07.95) EP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant: GIST-BROCADES B.V. [NL/NL]; Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL). (72) Inventors: BARENDSE, Rudolf, Carolus, Maria; Van Bosses-straat 9, NL-2613 CM Delft (NL). HARZ, Hans-Peter; 13 Mönchsbusch 22, D-67373 Dudenhofen (DE). BETZ, Roland, Jürgen; Im Brühl 5 205, D-67150 Niederkirchen (DE). (74) Agents: KLEIN, Bart et al.; Gist-Brocades N.V., Patents and Trademarks Dept., Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL).		(81) Designated States: AL, AM, AT, AZ, BB, BG, BR, BY, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: SALT-STABILIZED ENZYME PREPARATIONS (57) Abstract <p>The present invention discloses methods to improve the processing and storage stability of dry enzyme preparations. To this extent an inorganic salt, e.g. MgSO₄, is dissolved in an enzyme containing solution which is subsequently dried, using e.g. spray drying. Not only does the addition of salt improve the yield of enzyme activity upon drying, but also the storage and processing stability of the obtained solid composition containing the enzyme and the inorganic salt are improved with respect to enzyme activity. The invention further discloses solid compositions comprising enzymes and inorganic salts.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Salt-stabilized enzyme preparations

5

Field of the invention

The present invention relates to the conversion of enzyme solutions to
10 solid enzyme preparations with improved drying yield, and storage and
processing stability.

Background of the invention

15

The width of the application field and therefore the importance of enzymes
in modern technology is increasing rapidly. One of the major drawbacks of these
compounds however, is their aptitude to deactivation, caused not only by
extremes of temperature, pH and the like during processing, but also occurring
20 spontaneously on prolonged storage under otherwise normal conditions.

Often applied methods known in the art to diminish this problem include
the use of a variety of additives claimed to stabilize enzymes in solution or the
conversion of the enzyme solution to a dry formulation by means of freeze
drying, spray drying or other techniques suitable for this purpose. The conversion
25 of an enzyme in solution to a dry form is often obligatory when the application
so demands (e.g. convenient mixing with other dry components).

Although drying in itself is a valuable tool in the improvement of the
enzyme storage stability, the process step itself often causes a substantial loss
of activity and the final product is still susceptible to inactivation. This activity
30 loss during storage or processing is strongly dependent on moisture content of
the preparation and this therefore has to be most stringently controlled to
maintain the so valued product stability. This also includes a severe reduction of
choice in the compounds suited for addition to the final enzyme product for the
sake of standardization or other purposes.

35 In a variety of cases the dry enzyme is intended for applications in which
the enzyme has to be incorporated in a product in which the moisture content
can not be so strictly controlled. In these cases the enzyme stability is then
severely reduced.

40 Several inventions have been made in the field of stabilizing enzymes
against losses during drying and subsequent storage and handling.

The bulk of these inventions (such as presented in the patent (applications) US 3515642, EP 0501375A1 and WO 91/14773) are concerned with the addition of carbohydrate and, more specific, sugar or polyol components to the enzyme concentrate.

5 Also known in the art is the inclusion of components into the formulation with the aim to produce a glassy product at storage temperature, thus improving enzyme stability (EP 0383569A2).

Another approach is the addition to the formulation of one or several components able to bind moisture. This will reduce the water activity of the final preparation or temporarily prevent the interaction of water penetrating from the surroundings with the enzyme itself.

10 The use of organic and inorganic salts as a processing aid (e.g. to improve flowing behaviour of the product) or bulking/standardizing agent is well known. However, the use of inorganic salts to improve stability of dry enzyme preparations during processing or storage does not appear to be described in the art. In some cases enzyme destabilisation due to salt addition was even specifically mentioned (see e.g. Mikhailova et al. 1989, Vestsi Akad Navuk BSSR, Ser Biyal Navuk 6: 62-65).

15 EP 0522269A2 discloses the addition of insoluble calcium carbonate to an enzyme solution to be spraydried.

20 WO 92/11347 discloses the incorporation of water soluble inorganic salts into enzyme containing granulates. The water soluble inorganic salts must be chosen such that they do not affect the storage stability of the enzyme granulate. WO 92/11347 provides several sodium- and potassium-salts as being suitable for this purpose. The water soluble inorganic salts are however added to the enzymes in dry form prior to their granulation by extrusion.

30 Description of the invention

The present invention provides a method for the production of solid enzyme formulations with improved drying yield and improved storage and processing stability of the resulting products. This is achieved by preparing a solution comprising an enzyme and a water soluble inorganic salt, preferably an inorganic salt of a divalent cation, more preferably zinc, magnesium or calcium salt, most preferably zinc or magnesium sulphate, and subsequent drying of the product. Also a combination of salts as well as a combination of enzymes can

- 3 -

be used for this purpose. Divalent cations are preferred because they provide the best storage and processing stability. Sulphate is preferred as anion because it provides the best drying yield. The invention also provides solid enzyme formulations obtained by preparing a solution comprising an enzyme and a water soluble inorganic salt, preferably an inorganic salt of a divalent cation, more preferably zinc, magnesium or calcium salt, most preferably zinc or magnesium sulphate, and subsequent drying of the solution comprising the enzyme and the water soluble inorganic salt.

In the method of the invention, the salt is present while the enzyme is still in solution, i.e. prior to drying. Not only does this results in a higher yield during drying, but also the storage stability of the obtained dry enzyme preparations is improved as well as their processing stability. Processing stability is herein understood to mean the stability of the enzyme preparation during any handling of the enzyme preparation other than storage, such as e.g. the mixing of the enzyme preparation with other components or during the application of the enzyme.

Drying of the solution containing the enzyme and the salt will result in a solid composition which is homogeneous with respect to the distribution of the enzyme and the salt.

The drying of the enzyme and salt containing solution can be achieved by any drying method available to the skilled person, such as spray-drying, freeze drying, vacuum drying, fluid bed drying, and microwave drying. Drying of the enzyme-salt solution can also be combined with granulation methods which comprise e.g. the use of a fluid bed or a Multi-stage dryer (MSD). In case of the use of these granulation methods, the skilled person will understand that the obtained composition is not necessarily completely homogeneous. The obtained particles will usually consist of agglomerates of homogeneous particles or will consist of coated particles with a homogeneous core, with a homogeneous coating and/or combinations thereof. In any case there will be a homogenous interdispersion of the enzyme and the salt with respect to each other.

The specific examples of the invention demonstrate that the stabilising effect of the salt increases with increasing dosage of the salt to the enzyme solution, until at a certain point further increases in salt dosage no longer produce further improvement of the enzyme stability. For this reason at least 5%,

preferably at least 15%, more preferably at least 30%, still more preferably at least 60%, and most preferably at least 90% (w/w) of salt is added to the enzyme solution, wherein the salt dosage is expressed as the weight percentage (w/w) of added salt based on the weight of the enzyme in solution, not including
5 the weight of the crystal water of the salt crystals.

Although there is no upper limit to the addition of the salt from the view of the invention, a physical limit results from the maximum solubility of the salt in the enzyme concentrate. Higher salt/enzyme ratios can then only be realized by dilution of the enzyme solution or the use of a combination of salts.

10 Another point of consideration might be the dosage at which the enzyme is 'salted out', this being dependent on both the type of salt and the specific enzyme under consideration. Combinations of salts and enzymes giving rise to the 'salting out'-phenomenon are not excluded from the invention.

15 The invention can be used with enzymes or mixtures of enzymes from any type, including but not limited to phytases and other phosphatases, amylases, proteases, lipases and phospholipases, cellulases such as β -glucanases, pectinases, hemicellulases such as xylanases and other plant-cell wall degrading enzymes, esterases, rennets such as fungal proteases or chymosin, and β -galactosidases. It is to be understood that whenever referred to the enzyme or
20 an enzyme, also mixtures of enzymes are included in these terms, irrespective of whether such mixtures are obtainable directly in a single fermentation or by mixing enzymes obtainable in different fermentations; and further including enzymes obtainable by fermentation of recombinant organisms.

25 The enzyme is preferably selected from the group comprising fungal phytases, fungal hemicellulases, fungal cellulases and bacterial proteases. More preferably the enzyme is selected from the group comprising phytases and endo-xylanases derivable from a fungus which belongs to the *Aspergillus niger* Group as defined by Raper and Fennell (1965, In: The Genus *Aspergillus*, The Williams & Wilkins
30 Company, Baltimore, pp 293-344), β -glucanases and endo-xylanases derivable from a *Trichoderma* species, and proteases derivable from *Bacillus* species.

The invention also discloses solid compositions comprising at least one enzyme and an inorganic salt, wherein the dosage of the inorganic salt is at least
35 5% (w/w) of the weight of the enzyme(s). Preferably the inorganic salt is a salt of a divalent cation, more preferably zinc, magnesium or calcium salt, most preferably zinc or magnesium sulphate. The dosage of the inorganic salt in the composition is at least 5% (w/w), preferably at least 15%, more preferably at

- 5 -

least 30%, still more preferably at least 60% and most preferably at least 90% of the weight of the enzyme(s). Enzymes which may be prepared in compositions according to the invention include all those mentioned above, particularly those which are preferred.

5 In a further embodiment of the invention the above described solid compositions are used to prepare an animal feed. The preparation of animal feed often includes a pelleting step during which a significant amount of enzyme activity can be lost. The use of the solid composition of the invention reduces these losses in enzyme activity during pelleting. Typically an animal feed
10 composition comprises raw materials of vegetable origin providing energy and metabolites for growth. The feed is supplemented with minerals and vitamins and it is common to add animal fat and animal proteins to the feed.

In yet another embodiment of the invention, a solid composition
15 comprising a phytase with a improved storage stability is disclosed. The storage stability of the phytase in this solid composition is such that upon storage during 8 weeks, preferably in a closed container, at 30°C, less than 35%, preferably less than 20%, more preferably less than 10% and most preferably less than 5% of the initial phytase activity is lost. The phytase compositions may comprise an
20 organic or inorganic physiologically acceptable carrier (e.g. grain-based carriers like wheat middlings, wheat flour, rice hulls, and so on, other organic carriers like tapioca, potato starch, cane sugar and so on, inorganic carriers like sodium chloride, potassium sulphate, silica and so on), in combination with phytase enzyme, possibly combined with other enzymes for functionalities' or
25 conveniences' sake, and agents to further improve the products' appearance or functionality with respect to colour, flowing behaviour, physical, chemical or microbial stability, smell, taste, dust formation and so on. The skilled person will understand that similar compositions can be made for feed-enzymes other than phytase.

30

The following examples are provide by way of illustration of the invention.

- 6 -

ExamplesExample 1

5

A fungal phytase concentrate (as produced by and obtainable from Gist-brocades) containing approximately 11 % of enzyme by weight was split into two portions. To one portion 240 g/l of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ was added. After dissolution of the salt, resulting in a clear solution, both solutions were spray dried using a

10 Buchi lab scale spray drier with inlet temperature of 130°C and outlet temperature of 85°C.

The resulting powders were mixed with wheat middlings in a weight ratio of 1:10 to neutralize any differences in water activity between the preparations. After equilibration this was confirmed by measurement of the water activities,

15 which both were found to be 0.45 at room temperature.

The final products were then stored at 30°C in closed jars and after storage the products were simultaneously analyzed for enzyme activity. After eight weeks of storage the activity loss of the preparation without salt addition amounted to 35 %, against only 5 % loss for the preparation with salt addition.

20

Example 2

Into separate portions of a concentrate of a Bacillus-derived alkaline

25 protease (as produced by and obtainable from Genencor International Inc.) containing approximately 12% of enzyme protein by weight, 0 and 85 g/l of $\text{MgSO}_4 \cdot \text{OH}_2\text{O}$ was dissolved. These concentrates were then coated onto sodium sulphate carrier using a Niro STREA-1 lab scale fluid bed coater. The activity losses for this process step were 28% and 14% respectively.

30 The resulting particles were mixed with a surplus of bleach-containing commercial detergent powder for the European market and incubated in open jars at 35°C and 55% relative humidity. After 6 weeks of storage the remaining enzyme activity of the preparations was 42% with salt addition and only 30% without.

35

- 7 -

Example 3

240 g/l of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ was dissolved into a portion of Trichoderma-derived enzyme concentrate (as produced by and obtainable from Gist-brocades) with a total dry matter content of 25% by weight, containing both a β -glucanase and an endo-xylanase enzyme. The resulting solution and the original concentrate were separately coated onto a sodium sulphate carrier in a Glatt WSG-60 fluid bed coater. The process activity losses were 5% and 10% for the β -glucanase and 10% and 25% for the endo-xylanase with and without salt-addition respectively.

The resulting particles were incubated at 30°C and 62% relative humidity. After 12 weeks of storage the activity losses for the β -glucanase were 10% and undetectable and for the endo-xylanase the activity losses were 10% and 25%, for each enzyme with and without salt addition, respectively.

The granulates were also incorporated in a feed mixture and subjected to a pilot scale pelleting process at a temperature of 70°C. The β -glucanase activity losses amounted to 46% and 59% and the endo-xylanase losses were 37% and 66% with and without salt addition respectively.

The feed mixture used in the pelleting process was composed of:

<u>Components</u>	<u>% (w/w)</u>
Maize	50.00
Peas (22.9% raw protein)	3.50
Defatted soy (45.5% raw protein)	28.00
Tapioca	2.38
Animal meal (58% raw protein)	3.60
Fish meal (72.7% raw protein)	1.00
Hydrolysed feather meal	1.00
Soy oil	1.75
Animal fat	3.50
Vitamin/mineral premix	3.15
Limestone	0.86
Monocalcium phosphate	0.96
Sodium chloride	0.30

- 8 -

Example 4

Different amounts of magnesium sulphate were dissolved in 100 ml of demineralized water after which 100 ml of a fungal phytase concentrate (as produced by and obtainable from Gist-brocades) containing app. 17% by weight of pure enzyme was added to each portion. After mixing the resulting solutions were spray-dried using a lab-scale Büchi 190 Mini Spray Dryer with inlet/outlet temperatures of 140 and 80°C respectively.

The resulting powders were mixed with wheat middlings in a ratio of 1:9. After equilibration water activity values at 35°C were measured for control sample and sample with highest salt dosage to exclude water activity effects. The mixtures were subsequently incubated in closed bottles at 35°C for 8 weeks, after which activity losses were measured.

The results are presented in the following table:

MgSO ₄ -dosage [g/g enzyme]	Mixture Water Activity [-]	Storage losses after 8 weeks at 35°C
0	0.292	52%
0.31	-	37%
0.61	-	26%
1.19	-	15%
1.84	-	15%
2.38	0.303	17%

Example 5

According to the same protocol as mentioned in example 4 different salts in a dosage of 80 mmol per 100 ml of phytase concentrate have been used to produce stabilized enzyme powders. Mixing with wheat middlings and stability testing took place analogously to example 4.

The results are presented in the following table:

- 9 -

Salt type	Water Activity [-]	Spray Drying losses	Storage losses after 8 weeks at 35°C
None	0.29	6%	52%
magnesium sulphate	-	7%	15%
magnesium chloride	0.30	26%	43%
magnesium nitrate	0.31	32%	27%
zinc sulphate	-	5%	9%
zinc chloride	-	48%	5%
calcium chloride	-	40%	18%
calcium nitrate	-	44%	13%
sodium sulphate	-	11%	51%
potassium sulphate	-	17%	36%
ammonium sulphate	-	6%	46%

Example 6

Using the protocol described in Example 5, three phytase containing solutions were prepared containing either no added salt, magnesium sulphate or zinc sulphate. These solutions were subsequently dried using freeze drying instead of spray drying. The freeze drying was accomplished by freezing the mixed enzyme/salt solutions in a flask by submerging the flask in liquid nitrogen, after which high vacuum was applied to remove the water. The resulting dry preparations were crushed in a mortar and mixed with wheat middlings prior to stability testing for 8 weeks at 35°C in closed bottles. The results are presented in the following Table.

Salt type	Water activity of mixture	Activity losses after 8 weeks at 35°C.
None	0.31	46%
magnesium sulphate	0.31	16%
zinc sulphate	-	3%

Claims

1. A method to improve the processing stability or storage stability of a solid enzyme preparation by drying a solution containing an enzyme and a water soluble inorganic salt, whereby the dosage of the inorganic salt is at least 5% (w/w) of the weight of the enzyme.
2. A method for preparing a storage stable and processing stable solid composition comprising an enzyme, whereby the method comprises the steps of:
 - a) preparing a solution comprising an enzyme and a water soluble inorganic salt, wherein the dosage of of the inorganic salt is at least 5% of the weight of the enzyme, and
 - b) drying the solution.
3. A method according to claims 1 or 2, wherein the inorganic salt comprises a divalent cation.
4. A method according to claims 3, wherein the divalent cation is selected from the group consisting of zinc, magnesium and calcium.
5. A method according to any one of claims 1-4, wherein the inorganic salt comprises a sulphate anion.
6. A method according to any one of claims 1-5, wherein the enzyme is selected from the group comprising a phytase preferably from *Aspergillus*, a protease preferably from *Bacillus*, a hemicellulase preferably from *Aspergillus*, a hemicellulase and a cellulase both preferably from *Trichoderma*.
7. A method according to any one of claims 1-6, wherein the drying of the solution containing the enzyme and the inorganic salt comprises the use of a dryer selected from the group consisting of a spray dryer, a fluid-bed dryer and a freeze dryer.
8. A solid composition comprising at least one enzyme and a water soluble inorganic salt, wherein the dosage of the inorganic salt is at least

- 11 -

5% (w/w) of the weight of the enzyme and wherein the inorganic salt comprises a divalent cation.

- 5 9. A solid composition obtainable by:
- a) preparing a solution comprising an enzyme and a water soluble inorganic salt, wherein the dosage of of the inorganic salt is at least 5% of the weight of the enzyme, and
 - b) drying the solution.
- 10 10. A solid composition according to claims 8 or 9, wherein the salt is as defined in any one of claims 3-5.
11. A solid composition according to any one of claims 8-10, wherein
15 the enzyme is as defined in claim 6.
12. A solid composition according to any one of claims 8-11, wherein the composition is homogeneous with respect to the distribution of the inorganic salt and the enzyme.
- 20 13. A solid composition according to claim 8-11, characterized in that the composition comprises particles which are homogeneous with respect to the distribution of the inorganic salt and the enzyme.
- 25 14. A solid composition according to claim 8-11, characterized in that the composition comprises a coating which is homogeneous with respect to the distribution of the inorganic salt and the enzyme.
- 30 15. A solid composition comprising a phytase with a storage stability which is characterized in that upon storage of the solid composition during 8 weeks at 30°C, less than 35% of the initial phytase activity is lost.
16. An animal feed comprising a solid composition as defined in any one of claims 8-15.

35

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 96/03286

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/96 A23L1/00 C12N9/16 C12N9/24 C12N9/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 522 269 (BASF AG) 13 January 1993 see example 2	1-16
X	WO,A,92 11347 (HENKEL KGAA) 9 July 1992 see page 6, paragraph 1; claims 1,8	1,9-12
X	EP,A,0 501 375 (KAO CORP) 2 September 1992 see examples 1-3	1-16
X	INDIAN J BIOCHEM BIOPHYS, 19 (6). 1982 (RECD. 1983). 399-402., XP000609629 SOHONI S S ET AL: "ISOLATION AND CHARACTERIZATION OF A PROTEASE FROM BACILLUS-SUBTILIS" see figure 2	9-16
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

21 November 1996

Date of mailing of the international search report

29. 11. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Espen, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP 96/03286

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0522269	13-01-93	DE-A- 4119281 CA-A- 2070155 JP-A- 5153977 US-A- 5318903	17-12-92 13-12-92 22-06-93 07-06-94
WO-A-9211347	09-07-92	DE-A- 4041752 AT-T- 121770 DE-D- 59105338 EP-A- 0564476 ES-T- 2071478	25-06-92 15-05-95 01-06-95 13-10-93 16-06-95
EP-A-0501375	02-09-92	JP-A- 4271785	28-09-92
WO-A-9109943	11-07-91	DE-D- 69004101 DE-T- 69004101 EP-A- 0506866 ES-T- 2060360 JP-T- 5502791	25-11-93 24-03-94 07-10-92 16-11-94 20-05-93